

# NOTES ON FRESH-WATER COPEPODA IN THE UNITED STATES NATIONAL MUSEUM.

---

By C. DWIGHT MARSH,

*Of the United States Department of Agriculture, Washington.*

---

The collections on which these notes are based have been in the United States National Museum for some years, and most of them were obtained in connection with the work of the United States Fish Commission. The Florida collections were made by Dr. W. C. Kendall, those from Sodus Bay by Mr. A. C. Weed; the Mississippi, Louisiana, and Axton, New York, material was obtained by Dr. B. W. Evermann, and the western collections by Doctor Evermann, Drs. Evermann and Meek, Meek and Alexander, and Messrs. Cox and Ulrich.

Following is a list of the localities in which copepods were found and the names of the species identified in each:

## COPEPODA FOUND IN PLANKTON COLLECTIONS RECEIVED FROM UNITED STATES NATIONAL MUSEUM IN 1911.

Woods Hole, Massachusetts, fresh-water pond:

*Diaptomus leptopus*, var. *piscinæ* Forbes.

*Cyclops viridis*, var. *americanus* Marsh.

Sodus Bay, New York:

Hotel dock, Leroys Island, to sandbar—

*Diaptomus reighardi* Marsh.

*Cyclops albidus* Jurine.

*Cyclops viridis*, var. *brevispinosus* Herrick.

Sand bar to south end of Nub Island—

*Diaptomus reighardi* Marsh.

*Cyclops viridis*, var. *brevispinosus* Herrick.

East side, Leroys Island—

*Cyclops albidus* Jurine.

*Cyclops serrulatus*, var. *elegans* Herrick.

*Cyclops leuckarti* Claus.

*Cyclops viridis*, var. *brevispinosus* Herrick.

*Diaptomus reighardi* Marsh.

South end of Nub Island to hotel dock, Leroys Island—

*Cyclops viridis*, var. *brevispinosus* Herrick.

*Cyclops albidus* Jurine.

*Diaptomus reighardi* Marsh.

## Axton, New York: —

## Rock Pond—

*Cyclops leuckarti* Claus.*Cyclops strenuus* Fischer.

## Pickerel Pond—

*Cyclops leuckarti* Claus.

## Lake George, Florida:

## South end—

*Eurytemora affinis* Poppe.*Cyclops prasinus* Fischer.*Cyclops*, immature, probably *leuckarti*.*Diaptomus*, immature.

## Four miles south of Draytons Island—

*Cyclops prasinus* Fischer.*Diaptomus*, immature, probably *dorsalis* Marsh.

## Middle of lake—

*Diaptomus dorsalis* Marsh.*Cyclops leuckarti* Claus.

## Lake Monroe, Florida:

## Middle—

*Diaptomus dorsalis* Marsh.*Cyclops leuckarti* Claus.

## Upper end—

*Diaptomus dorsalis* Marsh.*Cyclops leuckarti* Claus.

## Middle, between Sanford and Enterprise—

*Diaptomus dorsalis* Marsh.*Cyclops leuckarti* Claus.

## Just below railroad wharf—

*Cyclops serrulatus* Fischer.

## Little Lake George, Florida:

## Opposite Beecher Point—

*Eurytemora affinis* Poppe.*Diaptomus dorsalis* Marsh.*Cyclops prasinus* Fischer.*Cyclops leuckarti* Claus.

## No locality indicated—

*Eurytemora affinis* Poppe.*Diaptomus dorsalis* Marsh.*Cyclops leuckarti* Claus.*Cyclops serrulatus* Fischer.*Cyclops prasinus* Fischer.

## Station No. 3, at black beacon 65—

*Eurytemora affinis* Poppe.*Cyclops prasinus* Fischer.*Diaptomus dorsalis* Marsh.

## Locality illegible—

*Eurytemora affinis* Poppe.*Cyclops prasinus* Fischer.*Diaptomus dorsalis* Marsh.

## About middle—

*Cyclops prasinus* Fischer.

## Station No. 1, black beacon 67—

*Eurytemora affinis* Poppe.*Cyclops leuckarti* Claus.

## St. John's River, Florida:

## Opposite Palatka—

*Cyclops prasinus* Fischer.*Cyclops*, immature, probably *leuckarti*.*Diaptomus*, immature.

## Fort Gates—

*Eurytemora affinis* Poppe.*Cyclops prasinus* Fischer.*Diaptomus dorsalis* Marsh.

## Black Bayou, Mississippi:

*Eurytemora affinis* Poppe.*Cyclops albidus* Jurine.

## Flat Lake, Louisiana:

*Cyclops albidus* Jurine.*Eurytemora affinis* Poppe.

## San Marcos, Texas:

*Cyclops albidus* Jurine, immature.

## Saginaw River, Michigan:

*Cyclops serrulatus* Fischer.*Cyclops albidus* Jurine.

## Gambliis Lake, Idaho:

*Epischura nevadensis* Lilljeborg.*Cyclops serrulatus* Fischer.*Cyclops leuckarti* Claus.*Cyclops albidus* Jurine.

## Lake Pend Oreille, Idaho:

## July 6—

*Cyclops bicuspidatus* Claus.*Diaptomus ashlandi* Marsh.*Epischura nevadensis* Lilljeborg.

## Mountain Creek, June 27—

*Epischura nevadensis* Lilljeborg.*Diaptomus ashlandi* Marsh.

## Alturas Lake, Idaho:

## 8 p. m.—

*Diaptomus*, no mature males, probably *tyrelli* Poppe.*Cyclops viridis* Jurine.

## 6.30 p. m.—

*Diaptomus*, numerous, immature.*Cyclops viridis*, var. *americanus* Marsh.

## 8.10 a. m.—

*Diaptomus*, numerous, immature, probably *tyrelli* Poppe.*Cyclops viridis*, var. *americanus* Marsh.

## 7 p. m.—Same as above.

## 8 p. m.—Same as above.

## Lake Washington, Washington, east side:

*Diaptomus ashlandi* Marsh.*Epischura nevadensis* Lilljeborg.

## Tsiltcoos Lake, Oregon:

*Epischura nevadensis* Lilljeborg.

## Crater Lake, Oregon, towing from landing to island:

*Diaptomus*, one female, perhaps *franciscanus* Lilljeborg.

## Del Monte Lake, Monterey, California:

*Cyclops viridis*, var. *americanus* Marsh, 2 specimens.

Hocketts Lake, California (in Kern River region):

No copepods, but from other collections from Hocketts Lake were identified—

*Cyclops albidus* Jurine.

*Cyclops serrulatus*, var. *montanus* Brady.

*Diaptomus signicauda* Lilljeborg.

While most of the localities are new for the species, in the majority of cases no special significance is to be attached to the new facts of distribution. *Cyclops bicuspidatus*, *C. leuckarti*, *C. albidus*, *C. serrulatus*, and *C. prasinus* occur not only all over the United States, but are common in Europe and Asia, while some of these species extend their range to Africa without any change of structure. *C. viridis* in its varieties *brevispinosus* and *americanus* occurs everywhere in the United States. The localities for *Epischura nevadensis*, *Diaptomus ashlandi*, and *D. tyrelli* are not new. The localities of the other species, however, are of particular interest and justify the more extended statement following.

**DIAPTOMUS LEPTOPUS, var. PISCINÆ Forbes.**

This variety was originally described from collections made in Montana. It has also been found in Manitoba, Alberta, and Colorado. *Diaptomus leptopus* has seemed to be confined largely to the Mississippi Valley, although it has been reported from Massachusetts. *Diaptomus leptopus*, var. *piscinæ* has not before been reported from the eastern United States, so that its occurrence at Woods Hole is a matter of some interest.

**DIAPTOMUS DORSALIS Marsh.**

This species was first found in collections made by Prof. E. A. Birge in southern Louisiana, and up to the present time this has been the only locality for the species. Its occurrence, therefore, in great numbers in the collections made in Florida is of importance as extending the range of the species, because without doubt it will be found at intervening points. It was found in Little Lake George, Lake Monroe, and in the St. Johns River as far down as Palatka.

**DIAPTOMUS REIGHARDI Marsh.**

Specimens of this species were found in the collections made by Mr. A. C. Weed in Sodus Bay, New York, July 26 and August 9, 1909. These specimens are of much interest. *D. reighardi* was described from material collected from a lake on Beaver Island in Lake Michigan, and from Intermediate Lake, in the northern part of the southern peninsula of Michigan. It was later found in collections made by Dr. R. H. Ward in Crooked Lake, Michigan. Although extensive collections have been made in the other lakes of Michigan and in Lake Michigan, it has never been found in any other locality, and the supposition has been that it was a rather closely localized species. From its

occurrence in Sodus Bay it is assumed that it will eventually be found in bodies of water between Michigan and New York, and from the fact that Sodus Bay is connected with Lake Ontario it might be expected in that lake. The New York specimens correspond in every detail with those found in Michigan.

#### EURYTEMORA AFFINIS Poppe.

This species is found very abundantly in many places on the coast of Europe, in salt, brackish, and fresh waters. It was first reported in America by Herrick,<sup>1</sup> who says that it lives in the shallow bays and estuaries along the Gulf of Mexico. Foster,<sup>2</sup> 1904, gives definite localities near New Orleans, and Pearse,<sup>3</sup> 1906, records its occurrence on Nantucket Island, Massachusetts. Therefore its occurrence in great numbers, in the museum collections from the St. Johns River and Little Lake George, is no more than would be expected. Its occurrence in Flat Lake, Louisiana, and Black Bayou, Mississippi, however, is of a good deal of interest. Flat Lake is about 40 miles from the Gulf, while Black Bayou is more than 200 miles from the Gulf, and has an elevation of 144 feet. Inasmuch as *Eurytemora affinis* is commonly considered a salt-water form which is capable, after migration, of continuing its existence in brackish or fresh water, it is rather surprising to find it at a place so remote from salt water as Black Bayou.

It may be noted, in this connection, that Nordqvist<sup>4</sup> states that *Eurytemora lacustris* is found in lakes from 207 to 252 meters above the sea.

#### CYCLOPS STRENUUS Fischer.

*Cyclops strenuus* FISCHER, Bull. Soc. Imp. Moscou, vol. 24, 1851, pp. 419-425, pl. 9, figs. 12-22.—G. O. SARS, Forh. Vid. Selsk. Christiania, 1863, p. 236.—BRADY, Mon. Copepoda Brit. Isles, vol. 1, 1878, pp. 104-105, p. 119, figs. 1-7.—HERRICK, 12th Ann. Rept. Geol. Nat. Hist. Surv. Minnesota, 1884, p. 147.—DADAY, Math. és természettud. Közl. Vonatk. a haz. viszony., 1885, pp. 216-218.—VOSSELER, Jahresh Ver. vat. Nat. Württemberg, 22 Jahrg., 1886, p. 195, pl. 14, figs. 18-22.—RICHARD, Rev. Sci. Bourbonnais, vol. 1, 1888, pp. 61-62.—LANDE, Materyjaly do Fauny Skorupiakov Widlonogich Krolestwa Polskiego. Widlonogi Swobodnie Zyjace. I, Rodzina Cyclopy, 1890, pp. 53-55, pl. 21, figs. 156-163, 165.—SCHMEIL, Zeitschr. Naturw. Halle, vol. 64, 1891, p. 24.—BRADY, Nat. Hist. Trans. Northumberland, Durham, and Newcastle-upon-Tyne, vol. 11, 1891, p. 73, pl. 2, figs. 1-4.—RICHARD, Ann. Sci. Nat. Zool., vol. 21, 1891, pp. 227-228, pl. 6, fig. 18.—SCHMEIL, Deutschlands freilebende Süßwasser-Copepoden, Part 1. Cyclopidae, 1892, pp. 39-51, pl. 2, figs. 12-15.—LANDE, Mem. Soc. Zool., vol. 5, 1892, p. 161.—RICHARD, Rev. Biol. Nord de la France, vol. 5, 1893, p. 4.—SCHMEIL, Abh. Naturf. Ges. Halle, vol. 19, 1893-5, pp. 20-23.—HERRICK, Second Rept. State Zool. Minn., 1895, p. 99, pl. 23, figs. 12-13.—STEUER, Verh. Zool.-bot. Ges., Wien, 1897, p. 4.—MATILE, Bull.

<sup>1</sup> Twelfth Ann. Rept. Geol. and Nat. Hist. Surv. Minn., 1884, p. 182.

<sup>2</sup> Second Rep. Gulf Biol. Station, pp. 73-74.

<sup>3</sup> Amer. Naturalist, vol. 40, p. 242.

<sup>4</sup> Die Calaniden Finlands, 1888, p. 57.

- Soc. Natur. Moscou, 1897, No. 1, p. 122, pl. 2, fig. 6.—LILLJEBORG, Kongl. Svenska Vet.-Akad. Handl., vol. 35, 1901, p. 28, pl. 2, figs. 20-25.—GRÆTER, Rev. Suisse Zool., vol. 11, 1903, pp. 514-523, pl. 15, figs. 15 and 33.—VAN DOUWE, Die Süßwasserfauna Deutschlands, Heft. 2, 1909, pp. 24-25, figs. 69-71.
- Cyclops brevicaudatus* CLAUS, Das genus Cyclops, 1857, pp. 34-35, fig. 12; Die frei lebenden Copepoden, 1863, p. 100.—LUBBOCK, Trans. Linn. Soc. London, vol. 24, 1863, pp. 200-201.—FRIČ, Arch. Naturw. Land. Böhmen, vol. 2, 1872, Abth. 4, p. 221, fig. 15.—HOEK, Tijdsch. Ned. Dierk. Ver., vol. 3, 1876, pp. 15-17, pl. 2, figs. 1-9.
- Cyclops furcifer* CLAUS, Arch. f. Naturg., Jahrg. 23, vol. 1, 1857, p. 208-209, pl. 11, figs. 14-16.
- Cyclops scutifer* G. O. SARS, Forh. Vid. Selsk. Christiania, 1863, p. 237.—LILLJEBORG, Kongl. Svenska Vet.-Akad. Handl., vol. 35, 1901, p. 33, pl. 2, figs. 26-27.
- Cyclops abyssorum* G. O. SARS, Forh. Vid. Selsk. Christiania, 1863, p. 238.—BRADY, Nat. Hist. Trans. Northumberland, Durham, and Newcastle-upon-Tyne, vol. 11, 1891, p. 73, pl. 3, figs. 1-9.
- Cyclops clausii* LUBBOCK, Trans. Linn. Soc. London, vol. 24, 1863, pp. 201-202, pl. 31, figs. 12-14.
- Cyclops vicinus* ULJANIN, Schrift. Ges. Freunden Naturw. Moskau, vol. 11, 1875, pp. 30-31, pl. 10, figs. 1-7.—LANDE, Materyjaly do Fauny Skorupiakow Widlonogich Krolestwa Polskiego. Widlonogi Swobodnie Zyjace, I, Rodzina Cyclopy, 1890, pp. 47-49, pl. 16, figs. 33-41; pl. 21, fig. 164.—BRADY, Nat. Hist. Trans. Northumberland, Durham, and Newcastle-upon-Tyne, vol. 11, pt. 1, 1891, p. 12, pl. 1, figs. 6-9.—LILLJEBORG, Kongl. Svenska Vet.-Akad. Handl., vol. 35, 1901, p. 26, pl. 2, figs. 16-19.
- Cyclops fedtschenkoi* ULJANIN, Schrift. Ges. Freunden Naturw. Moskau, vol. 11, 1875, pp. 31-32, pl. 12, fig. 10.
- Cyclops pulchellus* BRADY, Mon. Copepoda Brit. Isles, vol. 1, 1878, pp. 107-108, pl. 17, figs. 1-3.
- Cyclops claudiopolitanus* DADAY, Math. és természettud. Közl. Vonatk. a haz. viszony, 1885, pp. 227-229, pl. 1, figs. 14-18.
- Cyclops hungaricus* DADAY, Math. és természettud. Közl. Vonatk. a haz. viszony, 1885, pp. 230-233, pl. 2, figs. 9-12.
- Cyclops bodamicus* VOSSELER, Jahresh. Ver. Vat. Nat. Württemberg, 22 Jahrg., 1886, p. 193, pl. 5, figs. 13-18.
- Cyclops lucidulus* VOSSELER, Jahresh. Ver. Vat. Nat. Württemberg, 22 Jahrg., 1886, p. 196, pl. 5, figs. 1-12.—THALLWITZ, Jahresb. ornith. Beobacht. Stat. Sachsen, vol. 5, 1890, p. 80.
- Cyclops quadricornis* SOSTARIC, Rad. jugoslav. akad., vol. 92, 1888, pp. 62-64, pl. 1, fig. 5.
- Cyclops strenuus*, var. *vicina* FRIČ and VÁVRA, Arch. Naturw. Land. Böhmen, vol. 9, 1893, pp. 57-58, figs. 44a and 44b.
- Cyclops miniatus* LILLJEBORG, Kong. Svenska Vet.-Akad. Handl., vol. 35, 1901, p. 24, pl. 2, figs. 13-15.
- Cyclops kolensis* LILLJEBORG, Kong. Svenska Vet.-Akad. Handl., vol. 35, 1901, p. 21, pl. 2, figs. 11-12.

Inasmuch as *Cyclops strenuus* has not before been found in America, a somewhat detailed description of the species is given.


The cephalothorax (fig. 1) is broadly oval, the posterior segments being much wider than in most of the species of *Cyclops*; the breadth is about one-half the length. The front is markedly produced. The

posterior corners of the first and second segments in the female are rounded, the third is either slightly produced backward or rounded, while the fourth and fifth are produced outward and backward in prominent points. The form of the fourth and fifth segments of the cephalothorax is a marked characteristic of this species. The fifth segment varies considerably in its form. Figures 2 and 3 show forms from specimens collected near Damascus. Specimens from Axton duplicated these forms. The fourth and fifth cephalothoracic segments of the male do not have the characteristic form of those found in the female.


The first abdominal segment about equals in length the remaining segments of the abdomen; the anterior end, while much narrower than the last cephalothoracic segment is nearly twice as wide as the posterior. The posterior margins of all the abdominal segments except the last are dentate; the last segment is armed with minute spines.



FIG. 1.—CYCLOPS STRENUUS, FEMALE.  $\times 36$ .




FIGS. 2-3.—CYCLOPS STRENUUS, LAST CEPHALOTHORACIC SEGMENT.  $\times 73$ .



FIGS. 4-5.—CYCLOPS STRENUUS. 4, FURCA OF SPECIMEN FROM SYRIA. 5, FURCA OF SPECIMEN FROM AXTON, NEW YORK.  $\times 148$ .

The branches of the furca are diverging, elongate, the length relatively to the rest of the abdomen being variable, but frequently equalling the last three abdominal segments. The inner margin is ciliate, and running the length of the dorsal surface, there is a somewhat irregular cuticular ridge; this ridge is not found in males or in immature specimens. The outer margin has an indentation at about one-third its length, and the lateral seta is set well toward the end. The terminal setae are rather short and weak. Figure 4 shows the furca of a specimen from Syria and is typical in its structure. Figure 5 is from an Axton specimen and differs from the type in that it is shorter, the lateral seta is set well back from the end of the segment, and the cuticular ridge of the dorsal surface is lacking; inasmuch as no egg-bearing females were found in the Axton collections, it is probable that the specimens were not quite mature.

The first antennæ are 17-segmented and reach about to the third cephalothoracic segment. The twelfth segment has a sensory club. The last three antennal segments bear a hyaline lamella with a somewhat irregular row of very minute spines. This structure is



FIGS. 7-8.—CYCLOPS STRENUUS. 7, TERMINAL SEGMENTS OF ANTENNA OF FEMALE FROM SYRIA. 8, TERMINAL SEGMENTS OF FEMALE FROM AXTON, NEW YORK.  $\times 292$ .





FIG. 6.—CYCLOPS STRENUUS. SECOND ANTENNA.  $\times 148$ .

shown in figures 7 and 8. Figure 7 is from a Syrian specimen and figure 8 is from a specimen collected at Axton, New York. These are drawn to the same scale and show the relative sizes from the two

localities. The second antenna, shown in figure 6, has no distinctive characteristics.

The spinous armature of the terminal segments of the exopodites of the swimming feet is represented by the formula 3, 4, 3, 3. This formula held true in all the specimens which have passed through the author's hands. Schmeil says that it may be also 2, 3, 3, 3, or 3, 3, 3, 3. Figures 9 and 10 show the structure of the terminal segments of the second and fourth feet.



FIGS. 9-10.—CYCLOPS STRENUUS. 9, TERMINAL SEGMENT OF EXOPODITE OF SECOND FOOT. 10, TERMINAL SEGMENT OF EXOPODITE OF FOURTH FOOT.  $\times 148$ .

The fifth foot is composed of two segments. The basal segment is commonly broader than long, but is somewhat variable in its relative dimensions; it bears at its outer distal angle a rather short plumose seta. The second segment is about twice as long as broad; it bears at its distal extremity a long plumose seta and about midway of its inner margin a stout serrate spine; distad of the spine the segment is about half as wide as the

basal portion. There is a group of small spines at the outer distal angle of this segment and at the base of the lateral spine. Figure 11 is the fifth foot of a Syrian specimen and figure 12 the same structure from an Axton specimen, drawn to the same scale.

The form of the *receptaculum seminis* is shown in figure 13. The form of the labrum is shown in figure 14. The second row of teeth is very distinct. Not enough specimens were examined to make sure that this, which has not been mentioned by other authors, is a constant structure, but it is interesting to note that Brady<sup>1</sup> in his figure of the labrum has a line where the author has found minute teeth; the row of teeth by a lower magnification appears as a line.

In all the specimens examined there was an absence of the customary seta on the basal segment of the fourth swimming foot. This apparently has not been noted by other authors.

Schmeil gives as the average length of females 1.5 to 2.5 mm.


The females of the Axton collections average 1.35 mm, the largest examined measuring 1.525 mm.

*Cyclops strenuus* is widely distributed in Europe, Asia, and northern Africa, but has not been reported before from America. It is distinctly a cold water form occurring most abundantly in the colder months.


Dr. B. W. Evermann collected it in Rock Pond, Axton, New York, on April 30. It seems strange that it has never been seen before, for it is probable that it is a form of wide distribution in America.

In the synonymy, for the most part, Schmeil has been followed. Granting the limits of variation as stated by him, the species published by Lilljeborg in 1901, *scutifer*, *vicinus*, *miniatus*, and *kolensis* should be considered as varieties, and it

would be a matter of personal opinion whether or not it is worth while to recognize them under distinct names.



FIGS. 11-12.—CYCLOPS STRENUUS. 11, FIFTH FOOT OF SPECIMEN FROM SYRIA. 12, FIFTH FOOT OF SPECIMEN FROM AXTON, NEW YORK.  $\times 292$ .



FIGS. 13-14.—CYCLOPS STRENUUS. 13, OUTLINE OF RECEPTACULUM SEMINIS.  $\times 148$ . 14, LABRUM.  $\times 292$ .

## REFERENCES.

BRADY, G. S.

1878. A monograph of the free and semi-parasitic copepoda of the British Isles. 3 vols. Ray Society, London.  
1891. A revision of the British species of fresh-water Cyclopidae and Calanidae. Nat. Hist. Trans. Northumberland, Durham, and Newcastle-upon-Tyne, vol. 11, pt. 1, pp. 68-120.

CLAUS, C.

- 1857a. Das Genus Cyclops und seine Einheimischen Arten. Inaugural dissertation, Marburg.  
1857b. Weitere Mittheilungen über die einheimischen Copepoden. Archiv f. Naturgesch., Jahrg. 23, vol. 1, pp. 205-210.  
1863. Die frei lebenden Copepoden mit besonderer Berücksichtigung der Fauna Deutschlands, der Nordsee und des Mittelmeeres. Leipzig.

DADAY, E. VON.

1885. Monographia Eucopodorum liberorum in Hungaria hucusque repertorum. Matematikai és természettudományi Közlemények Vonatkozólag a hazai viszonyokra, vol. 19, 1883.

DOUWE, C. VAN.

1909. Die Süßwasserfauna Deutschlands, eine Excursionsfauna, edited by Brauer. Heft II, Copepoda, Ostracoda, Malacostraca, by van Douwe, Neresheimer, Vávra and Keilhack. Jena.

FISCHER, S.

1851. Beiträge zur Kenntniss der in der Umgegend von St. Petersburg sich findenden Cyclopiden. Bull. Soc. Imp. Moscou, vol. 24, pp. 409-438.

FOSTER, E.

1904. Notes on the free-swimming copepods of the waters in the vicinity of the Gulf Biologic Station, Louisiana. Second Report Gulf Biologic Station, 1903, Bull. 2, May, 1904, pp. 69-79.

FRIČ, A.

1872. Die Krustenthiere Böhmens. Archiv der Naturwiss. Landesdurchforsch. von Böhmen, vol. 2, Abth. 4, pp. 201-269.

FRIČ, A., and VÁVRA, V.

1893. Untersuchungen über die Fauna der Gewässer Böhmens. Archiv der Naturwiss. Landesdurchforsch. von Böhmen, vol. 9, No. 2.

GRÆTER, A.

1903. Die Copepoden der Umgebung Basels. Rev. Suisse Zool., vol. 11, pp. 419-541.

HERRICK, C. L.

1884. A final report on the Crustacea of Minnesota included in the orders Cladocera and Copepoda. 12th Ann. Rep. Geol. and Nat. Hist. Sur. Minn., pp. 1-190.

HERRICK, C. L., and TURNER, C. H.

1895. Synopsis of the Entomostraca of Minnesota. Second Rep. State Zoologist of Minn., Geol. and Nat. Hist. Sur. Minn.

HOEK, P. P. C.

1876. De vrijlevende zoetwater-Copepoden der Nederlandsche Fauna. Tijdsch. d. Nederl. Dierkund. Vereenig, vol. 3, pp. 1-36.

LANDE, A.

1890. Materyjaly do Fauny Skorupiakow Widlonogich (Copepoda) Krolestwa Polskiego. Widlonogi Swobodnie Zyjace. I. Rodzina Cyclopy (Cyclopidae). Warsaw.  
1892. Quelques remarques sur les Cyclopidés. Mém. Soc. Zool., vol. 5, pp. 156-173.

## LILLJEBORG, W.

1901. Synopsis specierum huc usque in Suecia observatarum generis Cyclopis. Kongl. Svenska Vet.-Akad. Handl., vol. 35, No. 4.

## LUBBOCK, J.

1863. Notes on some new or little-known species of freshwater Entomostraca. Trans. Linn. Soc. London, vol. 24.

## MATILE, P.

1897. Contribution à la faune des Copépodes des environs de Moscou. Bull. Soc. Natur. Moscou, Année 1897, No. 1, pp. 113-139.

## NORDQVIST, O.

1888. Die Calaniden Finlands. Helsingfors.

## PEARSE, A. S.

1906. Fresh-water Copepoda of Massachusetts. Amer. Nat., vol. 40, pp. 241-251.

## RICHARD, J.

1888. Cladocères et copépodes non marins de la faune de France. Rev. Sci. du Bourbonnais, vol. 1, pp. 57-70.  
1891. Recherches sur le système glandulaire et sur le système nerveux des Copépodes libres d'eau douce, suivie d'une révision des espèces de ce groupe qui vivent en France. Ann. Sci. Nat. Zool., vol. 12, pp. 113-270.  
1893. Copépodes recueillis par M. le Dr. Th. Barrois en Égypte, en Syrie et en Palestine. Rev. Biol. du nord de la France, vol. 5.

## SARS, G. O.

1863. Oversigt af de indenlandske Ferksvandscopepoder. Forhandling i Videnskabs-Selskabet i Christiania, 1862, pp. 212-262.

## SCHMEIL, O.

1891. Beiträge zur Kenntniss der freilebenden Süsswasser-Copepoden Deutschlands mit besonderer Berücksichtigung der Cyclopiden. Zeitschr. f. Naturw. Halle, vol. 64, pp. 1-40.  
1892. Deutschlands freilebende Süsswasser-Copepoden, I Theil, Cyclopidae.  
1893. Copepoden des Rhätikon-Gebirges. Abhandlungen der Naturforschenden Gesellschaft zu Halle, vol. 19, pp. 1-40.

## SOSTARIC, D.

1888. Prilog poznavanju faune slatkovodnih korepnjaka hrvatske. rad. jugoslavenske akademije znanosti i umjetnosti, vol. 92.

## STEUER, A.

1897. Copepoden und Cladoceren des süßen Wassers aus der Umgebung von Triest. Verhandl. der k. k. zoologisch-botanischen Gesellschaft in Wien. Jahrg. 1897.

## THALLWITZ, J.

1891. 5 Jahresber d. Ornithol. Beobachtungsstat Sachsen, pp. 75-80.

## ULJANIN, W. N.

1875. Crustaceen von Turkestan. Schriften d. Gesellsch. v. Freunden der Naturwissensch. zu Moskau, vol. 11, pp. 22-41.

## VOSSELER, J.

1886. Die freilebenden Copepoden Württembergs und angrenzender Gegenden. Jahresh. des Vereins f. vaterl. Naturkunde in Württemberg, 42 Jahrgang, pp. 167-204.